

RESQCPR BIBLIOGRAPHY

STUDIES THAT INCLUDED AN IMPEDANCE THRESHOLD DEVICE DURING THE PERFORMANCE OF ACTIVE COMPRESSION DECOMPRESSION CPR

Clinical (Human) Studies

1. Debaty G, Labarere J, Frascone RJ, et al. Gasping during cardiopulmonary resuscitation is associated with a higher likelihood of one year survival after cardiac arrest. *Circulation*. 2016;134:A19029.
2. Debaty G, Aufderheide T, Swor R, et al. The paradoxical association between pulmonary edema and survival with favorable neurological function after cardiac arrest. *Circulation*. 2014;130:A248.
3. Raubenolt A, Mastenbrook J, Franklin K, Fales W. Comparison of standard CPR versus CPR with an intrathoracic pressure regulator versus active compression decompression CPR plus an impedance threshold device during out-of-hospital cardiac arrest. *Prehosp Emerg Care*. 2014;18(1):155.
4. Frascone RJ, Wayne MA, Swor RA, et al, Yannopoulos D, Lurie KJ, O'Neil BJ, Holcomb RG, Aufderheide TP. Treatment of non-traumatic out-of-hospital cardiac arrest with active compression decompression cardiopulmonary resuscitation plus an impedance threshold device. *Resuscitation*. 2013; 84:1214-1222.
5. Frascone RJ, Wayne M, Mahoney B, et al. Active compression decompression cardiopulmonary resuscitation and augmentation of negative intrathoracic pressure is neuro-protective in patients with an out-of-hospital cardiac arrest. *Prehosp Emerg Care*. 2012;16(1):178.
6. Yannopoulos D, Holcomb R, Frascone RJ, et al. Determinants of ventricular fibrillation incidence as first recorded rhythm during out-of-hospital cardiac arrest and association with long term neurological outcomes. Observations from a large randomized clinical study. *Circulation*. 2012:126:A296.
7. Aufderheide TP, Frascone RJ, Wayne MA, et al. Standard cardiopulmonary resuscitation versus active compression-decompression cardiopulmonary resuscitation with augmentation of negative intrathoracic pressure for out-of-hospital cardiac arrest: a randomized trial. *Lancet*. 2011;377:301-11.
8. Wayne M, Tupper D, Swor R, et al. Improvement of long-term neurological function after sudden cardiac death and resuscitation: impact of CPR method and post-resuscitation care. *Resuscitation*. 2011; 82;Suppl 1:S3.
9. Plaisance P, Soleil C, Lurie KG, et al. Use of an inspiratory impedance threshold device on a facemask and endotracheal tube to reduce intrathoracic pressures during the decompression phase of active compression-decompression cardiopulmonary resuscitation. *Crit Care Med*. 2005;33(5):990-994.
10. Plaisance P, Lurie KG, Vicaut E, et al. Evaluation of an impedance threshold device in patients receiving active compression-decompression cardiopulmonary resuscitation for out of hospital cardiac arrest. *Resuscitation*. 2004;61:265-271.
11. Wolcke BB, Mauer DK, Schoefmann MF, et al. Comparison of standard cardiopulmonary resuscitation versus the combination of active compression-decompression cardiopulmonary resuscitation and an inspiratory impedance threshold device for out-of-hospital cardiac arrest. *Circulation*. 2003;108(18):2201-2205.
12. Plaisance P, Lurie KG, Payen D. Inspiratory impedance during active compression-decompression cardiopulmonary resuscitation: a randomized evaluation in patients in cardiac arrest. *Circulation*. 2000;101(9):989-994.

Pre-Clinical (Animal) Studies

1. Niforopoulou P, Iacovidou N, Lelovas P, et al. Correlation of impedance threshold device use during cardiopulmonary resuscitation with post-cardiac arrest acute kidney injury. *Am J Emerg Med.* 2017;10.1016/j.ajem.2017.01.040.
2. Debaty G, Lurie KG, Metzger A, et al. Reperfusion injury protection during basic life support improves circulation and survival outcomes in a porcine model of prolonged cardiac arrest. *Resuscitation.* 2016;105:29-35.
3. Metzger A, Lick M, Berger, P, et al. Active intrathoracic pressure regulation minimizes the need for vasopressors during post cardiac arrest care and improves cerebral hemodynamics. *Circulation.* 2016;134:A1776.
4. Ryu JJ, Moore JC, Yannopoulos D, et al. The effect of head up cardiopulmonary resuscitation on cerebral and systemic hemodynamics. *Resuscitation.* 2016;102:29-34.
5. Debaty G, Shin SD, Metzger A, et al. Tilting for perfusion: head-up position during cardiopulmonary resuscitation improves brain flow in a porcine model of cardiac arrest. *Resuscitation.* 2015;87:38-43.
6. Kwon Y, Debaty G, Puertas L, et al. Effect of regulating airway pressure on intrathoracic pressure and vital organ perfusion pressure during cardiopulmonary resuscitation: a non-randomized interventional cross-over study. *Scand J Trauma Resusc Emerg Med.* 2015;23:83.
7. Bartos JA, Matsuura TR, Sarraf M, et al. Bundled postconditioning therapies improves hemodynamics and neurologic recover after 17 min of untreated cardiac arrest. *Resuscitation.* 2015;87:7-13.
8. Debaty G, Metzger A, Rees J, et al. Enhanced perfusion during advanced life support improves survival with favorable neurological function in a porcine model of refractory cardiac arrest. *Crit Care Med.* 2015;45(5):1087-1095.
9. Debaty G, Matsuura TR, Bartos JA, et al. Sodium nitroprusside-enhanced cardiopulmonary resuscitation facilitates intra-arrest therapeutic hypothermia in a porcine model of prolonged ventricular fibrillation. *Crit Care Med.* 2015;43(4):849-855.
10. Sideris G, Magkoutis N, Sharma A, et al. Early coronary revascularization improves 24 hour survival and neurological function after ischemic cardiac arrest. a randomized animal study. *Resuscitation.* 2014;85(2):292-298.
11. Rees J, Puertas L, Metzger A, et al. The product of ET_{CO2} and an EEG measurement correlates with cerebral perfusion pressure during cardiopulmonary resuscitation in a swine model of cardiac arrest. *Prehosp Emerg Care.* 2014;18(1):155.
12. Burnett A, Salzman JG, Segal N, et al. Potential negative effects of epinephrine on carotid blood flow and ET_{CO2} during active compression-decompression CPR utilizing an impedance threshold device. *Resuscitation.* 2012;83(8):1021-1024.
13. Metzger A, Herman J, McKnite S, et al. Improved cerebral perfusion pressures and 24-hr neurological survival in a porcine model of cardiac arrest with active compression-decompression cardiopulmonary resuscitation and augmentation of negative intrathoracic pressure. *Crit Care Med.* 2012;40(6):1851-1856.
14. Pantazopoulos IN, Xanthos TT, Vlachos I, et al. Use of the impedance threshold devices improves survival rate and neurological outcome in a swine model of asphyxial cardiac arrest. *Crit Care Med.* 2012;40(3):861-868.
15. Schultz J, Segal N, Kolbeck J, et al. Sodium nitroprusside enhanced cardiopulmonary resuscitation (SNPeCPR) improves vital organ perfusion pressures and carotid blood flow in a porcine model of cardiac arrest. *Resuscitation.* 2012;83:374-377.
16. Yannopoulos D, Segal N, McKnite S, et al. Controlled pauses at the initiation of sodium nitroprusside-enhanced cardiopulmonary resuscitation facilitate neurological and cardiac recovery after 15 minutes of untreated ventricular fibrillation. *Crit Care Med.* 2012;40(5):1562-1569.
17. Metzger AK, Herman M, McKnite S, et al. Improved cerebral perfusion pressures and 24-hr neurological survival in a porcine model of cardiac arrest with active compression-decompression

- cardiopulmonary resuscitation and augmentation of negative intrathoracic pressure. *Crit Care Med.* 2012;40(6):1851-1856.
18. Schultz J, Segal J, Kolbeck J, et al. Sodium nitroprusside enhanced cardiopulmonary resuscitation prevents post-resuscitation left ventricular dysfunction and improves 24-hour survival and neurological function in a porcine model of prolonged untreated ventricular fibrillation. *Resuscitation.* 2011;82S:S35-S40.
 19. Schultz J, Segal J, Caldwell E, et al. Sodium nitroprusside-enhanced cardiopulmonary resuscitation improves resuscitation rates after prolonged untreated cardiac arrest in two porcine models. *Crit Care Med.* 2011;39(12):2705-2710.
 20. Yannopoulos D, Matsuura T, Schultz J, et al. Sodium nitroprusside enhanced cardiopulmonary resuscitation improves survival with good neurological function in a porcine model of prolonged cardiac arrest. *Crit Care Med.* 2011;39(6):1269-1274.
 21. Yannopoulos D, Halperin HR, Lurie KG. Lower extremity counterpulsation during the decompression phase of CPR improves hemodynamics and provides continuous forward carotid blood flow. *Circulation.* 2007;116(16):II-485.
 22. Srinivasan V, Nadkarni VM, Yannopoulos D, et al. Rapid induction of cerebral hypothermia is enhanced with active compression decompression plus inspiratory impedance threshold device cardiopulmonary resuscitation in a porcine model of cardiac arrest. *J Am Coll Cardiol.* 2006;47(4):835-841.
 23. Lurie KG, Barnes TA, Zielinski TM, McKnite SH. Evaluation of a prototypic inspiratory impedance threshold valve designed to enhance the efficiency of cardiopulmonary resuscitation. *Respir Care.* 2003;48(1):52-57.
 24. Bahlmann L, Klaus S, Baumeier W, et al. Brain metabolism during cardiopulmonary resuscitation assessed with microdialysis. *Resuscitation.* 2003;59(2):255-260.
 25. Langhelle A, Stromme T, Sunde K, et al. Inspiratory impedance threshold valve during CPR. *Resuscitation.* 2002;52:39-48.
 26. Raedler C, Voelckel WG, Wenzel V, et al. Vasopressor response in a porcine model of hypothermic cardiac arrest is improved with active compression-decompression cardiopulmonary resuscitation using the inspiratory impedance threshold valve. *Anesth Analg.* 2002;95(6):1496-1502.
 27. Voelckel WG, Lurie KG, Sweeney M, et al. Effects of active compression-decompression cardiopulmonary resuscitation with the inspiratory threshold valve in a young porcine model of cardiac arrest. *Pediatr Res.* 2002;51(4):523-527.
 28. Voelckel WG, Lurie KG, Zielinski T, et al. The effects of positive end-expiratory pressure during active compression decompression cardiopulmonary resuscitation with the inspiratory threshold valve. *Anesth Analg.* 2001;92(4):967-974.
 29. Lurie KG, Ingbar D, Sweeney M, et al. Potential role of the vocal cords during cardiopulmonary resuscitation. *Circulation.* 1998;98(17):I-478.
 30. Lurie KG, Coffeen P, Shultz J, et al. Improving active compression-decompression cardiopulmonary resuscitation with an inspiratory impedance valve. *Circulation.* 1995;91(6):1629-1632.

Pre-clinical results are not necessarily representative of clinical outcomes. Improper use of the ResQCPR System could cause ineffective chest compressions and decompressions, leading to suboptimal circulation during CPR and possible serious injury to the patient. The ResQCPR System should only be used by personnel who have been trained in its use. The ResQPUMP should not be used in patients who have had a recent sternotomy as this may potentially cause serious injury. Improper positioning of the ResQPUMP suction cup may result in possible injury to the rib cage and/or internal organs, and may also result in suboptimal circulation during ACD-CPR.